



Anti-CD68 Mouse Monoclonal Antibody(5E2)

Catalog No.	EM42007-01	EM42007-02	
Size	50µl	100µl	
Species	Human Mouse Rat	Molecular	37KD

Background: The CD68 antigen is a 37kD transmembrane protein that is post-translationally glycosylated to give a protein of 87-115kD. CD68 is specifically expressed by tissue macrophages, Langerhans cells and at low levels by dendritic cells. It could play a role in phagocytic activities of tissue macrophages, both in intracellular lysosomal metabolism and extracellular cell-cell and cell-pathogen interactions. It binds to tissue- and organ-specific lectins or selectins, allowing homing of macrophage subsets to particular sites. Rapid recirculation of CD68 from endosomes and lysosomes to the plasma membrane may allow macrophages to crawl over selectin bearing substrates or other cells.

Specificity: The CD68 antibody detects endogenous CD68 proteins.

Application: IHC 1:200

Optimal dilutions/concentrations should be determined by the end user.


Form: Liquid ,1mg/ml

Storage buffer: PBS, pH 7.4, containing 0.02% **sodium azide** as Preservative and 50% Glycerol.

Storage: Store at -20°C. Do not aliquot the antibody.

EMAR

EMAR



IHC staining of human tonsil tissue CD68 mouse mAb diluted at 1:200.